

Ch. 2 - Measurement & Stats

$$S = \sqrt{\frac{\sum_{i=1}^N (x_i - \bar{x})^2}{N - 1}}$$

Measurement & Stats

- Why numbers?
- Distribution & Graphs : Histogram
- Central Tendency
- Mean, SoR, SSR, Variance, Standard Deviation
- In-class exercise
- Population vs. Sample
- Measurement Scales
- Precision vs. Accuracy
- Logic and Logical Fallacies Descriptive vs. Inferential Statistics
- Norms


Basic Statistics

- Why use numbers?
 - Pros:
 - convenient, succinct
 - universal
 - well-defined, repeatable
 - Cons:
 - precision vs. accuracy
 - numerical fallacy

Tabular Data

Year	Rainfall (inches)
1965	8.81
1966	14.76
1967	10.86
1968	7.86
1969	11.48

Data Distributions



Histogram


- Frequency Distribution
- Invented by Karl Pearson
- Shows data from *one* variable only
- Data is (often) collected into groups (“bins”)

280

Psychology 402 - Spring 2020 - Dr. Michael Diehr

Histogram

Frequency Distribution of San Diego Annual Rainfall 1965-2017



283

Psychology 402 - Spring 2020 - Dr. Michael Diehr

Describing Distributions

- Why? Large lists are inconvenient. Reduce many data points to a few numbers.
- Issue: Reducing data (“Degrees of freedom”) : throws away data.
- We are modeling our data using a simplification.
- “All models are wrong, some models are useful”
- Simple vs. Simplistic?

288

Psychology 402 - Spring 2020 - Dr. Michael Diehr

Descriptive Statistics

- Statistical Assumptions: When these are not met, weird things happen.
- Joe Smith is 6 feet tall, his child is 1 foot tall. Thus, the average height in the Smith household is 3.5 feet.
- If you are sitting in bar, and Bill Gates walks in, suddenly everyone in the bar is (on average) a multi-millionaire.

289

Psychology 402 - Spring 2020 - Dr. Michael Diehr

Alternative Notation

- Square Root (x) \sqrt{x} $\sqrt[2]{x}$
- X-Squared X^2 X^{**2} X^2
- Sum(x) $x_1+x_2+x_3\dots$ $\sum_{i=1}^N x_i$ $\sum x$
- Mean M $\frac{\sum x}{N}$ \bar{X}

290

Psychology 402 - Spring 2020 - Dr. Michael Diehr


Central Tendencies

- Values tend to cluster around a point.
- **Mean** : most common statistic, commonly referred to as the “average”. Formula $\Sigma X / N$
- **Mode**: the most common value in a set
 - rare to use in statistics
- **Median**: the middle-most value in a set
 - the value at which half are above and half are below. Aka the 50th percentile.


291

Psychology 402 - Spring 2020 - Dr. Michael Diehr


Histogram



Normal Distribution




Skewed Distribution



Skew

- negative skew : fatter tail on the left
- positive skew : fatter tail on the right



Measures of Central Tendency 1

	Description	Algorithm	Formula
Mean	the “average”	sum values, divide by N	$\bar{x} = \frac{\sum_{i=1}^N x_i}{N}$
Median	the “middle-most” value	sort values, find middle value	50th percentile
Mode	the “most common” value	find most frequent value	...

Measures of Central Tendency 2

Behavior:	Normal Distribution	Skewed Distribution
Mean	same	overly affected by outliers
Median	same	fairly resistant to outliers
Mode	same	resistant to outliers

Measures of Dispersion 1

- Compare each measured value to the average
- “for a typical value, how far away is it from the mean”
- “Difference score” or “residual” can be calculated as the difference between the actual score and the mean. In other words, $d_i = x_i - \bar{X}$
- Take the average (mean) of the difference scores.
- Average difference score = $\text{Sum}(d) / N$

300

Psychology 402 - Spring 2020 - Dr. Michael Diehr

Average Difference Score

	Score (x)	Mean (\bar{X})	Difference $d = (x - \bar{X})$
	2	6	-4
	3	6	-3
	9	6	3
	11	6	5
	14	6	8
	1	6	-5
	6	6	0
	4	6	-2
	5	6	-1
	5	6	-1
Sum	60	60	0
Mean	6	6	0

301

Psychology 402 - Spring 2020 - Dr. Michael Diehr

Sum of Residuals

- Given N samples of x : $x_1, x_2, x_3 \dots x_N$
- mean of x $\bar{x} = \frac{\sum_{i=1}^N x_i}{N}$
- residuals $d_i = x_i - \bar{x}$
- Sum of Residuals is always zero

$$\sum_{i=1}^N d_i = 0$$

302

Psychology 402 - Spring 2020 - Dr. Michael Diehr

Sum of Residuals

- The “average difference score” score will *always* equal zero
- Solution:
 - Square the residuals *before* adding: removes the negative values.
 - “SSR” or Sum of Squared Residuals

303

Psychology 402 - Spring 2020 - Dr. Michael Diehr

SSR: Sum of Squared Residuals

- Given N samples of X: $x_1, x_2, x_3 \dots x_N$
- mean of x $\bar{x} = \frac{\sum_{i=1}^N x_i}{N}$
- residuals $d_i = x_i - \bar{x}$
- Sum of Squared Residuals (SSR)

$$SSR = \sum_{i=1}^N (d_i)^2$$

304

Psychology 402 - Spring 2020 - Dr. Michael Diehr

Problems with SSR

- SSR depends on units used (e.g. meters vs. millimeters)
- SSR depends on N (# of samples)
- Therefore, it's hard to interpret
 - is SSR = 0.00342 high or low?
 - is SSR = 2343249 high or low?

305

Psychology 402 - Spring 2020 - Dr. Michael Diehr

Remove the influence of N

- The Sum of a set of values depends on the number (N) of values:
- $\sum_{i=1}^N x_i$
- Take the average (mean) : divide by N, which removes the influence of N.

306

Psychology 402 - Spring 2020 - Dr. Michael Diehr

Variance

- Problem: SSR depends on N
- Solution: Take the average of SSR to remove the influence of N
- The average of the squared residuals is called Variance (S^2)

307

Psychology 402 - Spring 2020 - Dr. Michael Diehr

Variance

- Variance = SSR/N
- Variance = mean of squared residuals

$$S^2 = \frac{\sum_{i=1}^N (d_i)^2}{N}$$

$$S^2 = \frac{\sum_{i=1}^N (x_i - \bar{x})^2}{N}$$

308

Psychology 402 - Spring 2020 - Dr. Michael Diehr

Problems with Variance

- Units are squared:
 - measuring height in meters? variance is meters²
 - measuring # of cupcakes eaten? variance is (# of cupcakes eaten)²
- Won't someone rid me of these meddlesome squared units?

309

Psychology 402 - Spring 2020 - Dr. Michael Diehr

Standard Deviation

- Improving on Variance:
- The square root of Variance (S^2) gives S, which is called "Standard Deviation".
- Also abbreviated SD, StdDev or σ (Greek letter sigma)
- SD : easier to understand because it's in the same units as your measurement.
- SD is a unique property of the normal distribution -- given a mean and a SD you have uniquely specified the distribution

310

Psychology 402 - Spring 2020 - Dr. Michael Diehr

Standard Deviation

- SD = Square root of Variance


$$S = \sqrt{\frac{\sum_{i=1}^N (d_i)^2}{N}}$$

$$S = \sqrt{\frac{\sum_{i=1}^N (x_i - \bar{x})^2}{N}}$$

311

Psychology 402 - Spring 2020 - Dr. Michael Diehr

Normal Distribution




In a normal distribution, about 68.2% of values fall within ± 1 SD

312

Psychology 402 - Spring 2020 - Dr. Michael Diehr

SSR, Variance and SD



314

Psychology 402 - Spring 2020 - Dr. Michael Diehr

Central Limit Theorem

- No matter the shape of the Population distribution, if you take enough (*) samples of the mean, the distribution of your samples of the mean will have a Normal distribution
- Central Limit Theorem Exercise (Javascript)
- This fact makes our life easy: Many statistics assume a normal distribution. The CLT provides us a normal distribution in most cases, even when the population data is skewed

317

Psychology 402 - Spring 2020 - Dr. Michael Diehr

Exercise: normal distribution

- Roll one 10-sided die 10 times and record the results
- Prediction
 - Your Distribution: Uniform (flat)
 - Mean : 4.5
 - Class Distribution: ???
- hint: What is N? # die rolls, # of students?
- List and Graph results
- Does the distribution look normal?
 - if so, why?

319

Psychology 402 - Spring 2020 - Dr. Michael Diehr

Exercise 1: Die Rolls


X	M	$d = (x_i - M)$	$(\text{residual})^2$
2	3	-1	1
3	3		
5	3		
2	3		

N	$M = \bar{X}$	$\Sigma \text{Residuals}$	$\Sigma (\text{residual}^2)$	$\frac{\Sigma (\text{residual}^2)}{N-1}$	$\sqrt{S^2}$
	3				

320

Psychology 402 - Spring 2020 - Dr. Michael Diehr


Graph:



Caption: _____

322

Psychology 402 - Spring 2020 - Dr. Michael Diehr



Exercise: normal distribution 2

- Compute Mean (\bar{X}) - is it near 4.5?
- Compute residuals
- Compute sum of residuals -- do they add to zero?
- Compute squared residuals
- Compute Sum of squared residuals (SSR)
- Divide SSR by (N-1) - this is Variance or (S^2)
- Take square root of variance - this is S or Standard Deviation
- For this exercise, SD should be near 2.8

Psychology 402 - Spring 2020 - Dr. Michael Diehr

341

Review

- Themes
 - 18th-19th century
 - 19th-20th century
- Theories of Human Development
 - Creationism
 - Polygenism
 - Evolution
 - Genetics
- Controversy
 - IQ testing of various groups

355

Psychology 402 - Spring 2020 - Dr. Michael Diehr


Review

- Why Numbers?
 - pros
 - cons
- Distributions
 - Tables
 - Graphs

356

Psychology 402 - Spring 2020 - Dr. Michael Diehr

SSR, Variance and SD




- Frequency Distribution aka Histogram
 - graphically shows data
- Central Tendency
 - mean, median, mode
- Dispersion or Variation
 - residual
 - sum of residuals = 0
 - sum of squared residuals > 0
 - SSR/N = Variance
 - $\text{Sqrt}(\text{Variance}) = \text{Standard Deviation}$

Psychology 402 - Spring 2020 - Dr. Michael Diehr

358

Review

Review : Distributions



In a normal distribution, about 68.2% of values fall within ± 1 SD

359

Psychology 402 - Spring 2020 - Dr. Michael Diehr

Standard Deviation

- SD = Square root of Variance

$$S = \sqrt{\frac{\sum_{i=1}^N (d_i)^2}{N}}$$

$$S = \sqrt{\frac{\sum_{i=1}^N (x_i - \bar{x})^2}{N}}$$

Psychology 402 - Spring 2020 - Dr. Michael Diehr

360

Exercise: key points

- Some events (such as the roll of a die) have a flat (or ‘uniform’) distribution, but these are rare.
- Many big events are composed of many small events.
- Events in the real world often are distributed in a (nearly) normal distribution
- Assuming a normal distribution, the easiest way to describe the data is by two factors: Mean and SD.

361

Psychology 402 - Spring 2020 - Dr. Michael Diehr

Types of Statistics

- Descriptive:
 - Goal: help you describe the data
 - reduce the amount of data necessary for understanding
 - don’t draw conclusions -- “just the facts”
- Inferential:
 - Goal: draw conclusions from your sample to the larger data set (population)

361

Psychology 402 - Spring 2020 - Dr. Michael Diehr

Measurement Scales

- Nominal
- Ordinal
- Interval
- Ratio

382

Psychology 402 - Spring 2020 - Dr. Michael Diehr

Nominal Scale

- Nominal: Name or ID only
 - red, blue, green....
 - john, tony, fred...
 - Sci2-243, Sci2-245...
- does not signify Ordering, Ranking, or More/Less
- Gotcha: even if used with Numbers it may be still a Nominal.
- Example: colors, names, room numbers, ID numbers

383

Psychology 402 - Spring 2020 - Dr. Michael Diehr

Ordinal Scale

- Ordinal : ordering
 - first, second, third....
1, 2, 3...
A, B, C...
 - signifies Order, but can't assume distance between items is the same, e.g. the difference between an A and a B may be much different than a B and a C
- Example: Class Rank, Assignment Grade, Product Ratings

384

Psychology 402 - Spring 2020 - Dr. Michael Diehr

Interval Scale

- Interval: specifies orders AND inter-item distance
 - -3, -2, -1, 0, 1, 2, 3.... 100, 105, 115
 - the difference between two numbers IS the same, e.g. 100 to 105 should be the same amount as 105 to 110
- Does NOT have an absolute zero.
- Example: temperature in Degrees Farenheit

385

Psychology 402 - Spring 2020 - Dr. Michael Diehr

Ratio Scale

- Ratio: specifies orders AND inter-item distance and has absolute zero
 - 0, 1, 2, 3.... 100, 105, 115
 - the difference between two numbers IS the same, e.g. 100 to 105 should be the same amount as 105 to 110
- Does have an absolute zero.
- Example: temperature in Degrees Kelvin

386

Psychology 402 - Spring 2020 - Dr. Michael Diehr

Measurement Scales

	Magnitude	Equal Intervals	Absolute Zero
Nominal			
Ordinal	✓		
Interval	✓	✓	
Ratio	✓	✓	✓

387

Psychology 402 - Spring 2020 - Dr. Michael Diehr

Scales: Practical Info

- Nominal Scale: common
 - common stats: Count, Frequency, Mode
- Ordinal Scale: less common
 - stats: specialized “nonparametric” techniques required
- Ratio and Interval: common
 - Often can be treated identically with same statistical techniques

388

Psychology 402 - Spring 2020 - Dr. Michael Diehr

Descriptive Statistics

- Count (N)
- Range (minimum, maximum)
- Frequency Distribution (histogram)
- Rank order, percentile (%ile)
- Central Tendency
 - Mean
 - Median
 - Mode
- Variation / Dispersion (Variance, Standard Deviation)

389

Psychology 402 - Spring 2020 - Dr. Michael Diehr

Population vs. Sample

- Ideally, measure *everyone* to get the exact value (*Population parameter*)
- Practically, this is impossible.
- Take samples instead, and calculate the *Sample statistic*.
- The “Law of Large Numbers”, “Sampling Theory”, “Central Limit Theorem” makes life easier
- [Central Limit Theorem Exercise \(Javascript\)](#)
- Some formulas differ for *Population* vs. *Sample* (divide by N or divide by N-1 ?)

399

Psychology 402 - Spring 2020 - Dr. Michael Diehr

Population v. Sample

	Population	Sample
Definition	the entire set of items	the actual subset you measured
Descriptives	“Parameters”	“Statistics”
Symbols	Greek	Roman
Mean	μ	\bar{x}
Std. Deviation	σ	s
Variance	σ^2	s^2
Divide by	N	N-1

400

Psychology 402 - Spring 2020 - Dr. Michael Diehr

Law of Large Numbers

- If you take enough* samples, the sample mean approaches the population mean.
- Example: a coin has two sides. If heads=1 and tails = 0, then the average expected result is exactly 50% Heads (0.5) in the long run.
- However, if you flip a coin just a few times, getting exactly 0.5 is not likely.
- The LLN states that you will if you take enough samples.

* what is “enough”? Rule of thumb : 100.

404

Psychology 402 - Spring 2020 - Dr. Michael Diehr

LLN Demonstration

- Law of Large Numbers
- [Demonstration with Coin Flips](#)

405

Psychology 402 - Spring 2020 - Dr. Michael Diehr

Logical Fallacies

406

Psychology 402 - Spring 2020 - Dr. Michael Diehr

Logical Arguments

- Logical arguments or inferences generally have several components:
 - Premises
 - Conclusions
- Example:
 - Premise: All English people are musicians
 - Premise: John Lennon was English
 - Conclusion: John Lennon was a musician

408

Psychology 402 - Spring 2020 - Dr. Michael Diehr

Logical Arguments 2

- An Inference can be either Valid or Invalid -- this refers to the Structure of the argument (not the Facts themselves)
 - All A are B
 - All C are A
 - All C are B
- A Valid inference can still come to a false conclusion, and vice-versa

409

Psychology 402 - Spring 2020 - Dr. Michael Diehr


Logical Fallacies

- A Logical Fallacy generally means that your inference is Invalid to begin with. In addition, your facts may or may not be true, but the flaw in reasoning has occurred before you even apply facts.
- Example: Affirming the consequent
- If P, then Q bank owners are rich
Q is true Bill Gates is rich
Therefore P Bill Gates works at a bank

410

Psychology 402 - Spring 2020 - Dr. Michael Diehr

Cognitive Bias Codex



411

Dr. Michael Diehr

Psychology 402 - Spring 2020 - Dr. Michael Diehr

Biased Sample

- Every individual x that we have seen from sample X has characteristic Z
Therefore ALL X have characteristic Z
- Every student I talk to in this class is interested in Psychology
Therefore, ALL students are interested in Psychology

412

Nominal and Numeric

- Nominal Fallacy:** The tendency to believe that something has a name or identification, it exists or has special meaning.
“I am sleepy” vs. “I am suffering from activity-induced-rest-reduction-performance-impairment syndrome”
- Numerical Fallacy:** belief that something has been measured and assigned a number, it actually exists. “I’m really sad” vs. “I scored a 32 on the Beck Depression Inventory”

413

Psychology 402 - Spring 2020 - Dr. Michael Diehr

Reification Fallacy

- To Reify - to make something more concrete or real
- Examples:
 - “An A student”
 - “High IQ”
 - “Top of the class”
 - “A F Grade”

414

Psychology 402 - Spring 2020 - Dr. Michael Diehr


Ranking Fallacy

- Reducing a complex phenomenon (e.g. intelligence), giving it a single number (reification) and then ordering based on that number
- Examples:
 - A IQ of 93 is better than an IQ of 90
 - An income of \$50,000 is better than \$45,000

415

Psychology 402 - Spring 2020 - Dr. Michael Diehr

Hasty Generalization



417

Psychology 402 - Spring 2020 - Dr. Michael Diehr

Other Fallacies

- Begging the question -- circular argument
- Correlation implies causation
- Post hoc ergo propter hoc (*after this, therefore because of this*)
- Appeal to Authority
- Ad-hominem
- Straw Man
- False Dilemma

418

Psychology 402 - Spring 2020 - Dr. Michael Diehr

Significant Figures

- “digits of precision” or “sig. fig.”
- Ignoring *leading zeros*, how many digits in the measurement?
 - 00123 has 3 sig. fig.
 - 12003 has 5 sig fig
 - -9.87 has 3 sig fig
 - 0.000987 has 3 sig. fig.
 - 12.1 has 3 sig. fig.
 - 12.0 has 3 sig. fig.
- Please use 3 sig fig for this class
- Note: not the same as “decimal places”

419

Psychology 402 - Spring 2020 - Dr. Michael Diehr

Precision vs. Accuracy

- Precision : the level of detail a measurement is made with, often specified with an error-range
 - “about 6 feet plus or minus 1 foot” vs. “6 foot 11 inches plus or minus 1 inch”
- Accuracy: how close the measured value is to the actual value, does it “hit the target”
 - Think arrow vs. shotgun
- A number can be precise and accurate, precise but inaccurate, or accurate but imprecise.

420

Psychology 402 - Spring 2020 - Dr. Michael Diehr


Precision Fallacy

- A number that is *precise* may seem to be *accurate* when it is not
- A measurement that is *reliable* may seem to have *validity* when it does not

421

Psychology 402 - Spring 2020 - Dr. Michael Diehr

Precision vs. Accuracy



- Target shooting analogy
- Similar to Reliability vs. Validity

Psychology 402 - Spring 2020 - Dr. Michael Diehr

422

423

Psychology 402 - Spring 2020 - Dr. Michael Diehr

Fallacies re: Probability

- Classical
- Gambler's Fallacy
- Bayesian Reasoning

9 Heads in a row

- You are flipping a coin, and get 9 heads in a row
H H H H H H H H H
- What is the % chance the next flip will be a H ?
- Three common answers:
 - 50/50
 - more likely Heads
 - more likely Tails

424

Psychology 402 - Spring 2020 - Dr. Michael Diehr

425

Psychology 402 - Spring 2020 - Dr. Michael Diehr

9 Heads: Classical Inference

- Coin flips are independent 50/50 events, therefore 50% : Logical/Statistical
- This is the **correct** answer for a fair coin

LLN Demonstration

- Law of Large Numbers
- Demonstration with Coin Flips

426

Psychology 402 - Spring 2020 - Dr. Michael Diehr

427

Psychology 402 - Spring 2020 - Dr. Michael Diehr

9 Heads: Gambler's Fallacy

- Coin flips are independent 50/50 events, but we just saw 9/10 heads, therefore a Tail is "due"
- This is the "Gambler's Fallacy" and one reason Casinos make tons of money. The reasoning is faulty.
- Note: when dealing with draws w/o replacement, this logic is **correct**. For example, a single-card blackjack deck -- if no face cards have come up after 30 cards, then face cards are due

9 Heads: Bayesian Statistics

- Coin flips are supposed to be 50/50 events, but we just saw 9/10 heads, therefore the data is telling us that perhaps this is not a fair coin.
- Bayes' theorem suggests you evaluate the prior probabilities in determining future behavior
- In this case, you'd conclude that Head is more likely

428

Psychology 402 - Spring 2020 - Dr. Michael Diehr

429

Psychology 402 - Spring 2020 - Dr. Michael Diehr

Louis Agassiz

- Swiss-born, European-trained biologist / geologist
- Came to Harvard in 1847
- Creationist -> Polygenist
- Taxonomist
- Resisted Darwin's theory of Evolution
- d. 1873




Psychology 402 - Spring 2020 - Dr. Michael Diehr

460

Review: Samuel George Morton


- Theory of Polygenism
 - Humans are composed of different species, created by god
- Craniometry
- Biological Determinism
- “Scientific Racism”
- The “American School”
- d. 1851



Psychology 402 - Spring 2020 - Dr. Michael Diehr

461


Crania Americana



Samuel George Morton
1839

462

Psychology 402 - Spring 2020 - Dr. Michael Diehr



Psychology 402 - Spring 2019 - Dr. Michael Diehr

464

Morton's Data as printed

Race	N	Cranial Volume Mean
Caucasian	52	87
Mongolian	10	83
American	144	82
Malay	18	81
Ethiopian	29	78

466

Psychology 402 - Spring 2020 - Dr. Michael Diehr

Data, corrected

Race	Mean (Morton)	Mean (corrected)
Caucasian	87	87
Mongolian	83	87
American	82	86
Malay	81	85
Ethiopian	78	83

467

Psychology 402 - Spring 2020 - Dr. Michael Diehr

Seed vs. Shot

Race	Difference (seed - shot)
Caucasian	1.8
Mongolian	n/a
American	2.2
Malay	n/a
Ethiopian	5.4

469

Psychology 402 - Spring 2020 - Dr. Michael Diehr

Morton's errors

- Fundamental arithmetic errors
- Data selection errors
- Failure to measure or control for external variables (biological sex, body size, etc.)
- Basic Statistical errors (averaging measurements from unequal size subgroups)
- The racist thumb press?
- Is he a liar? Conscious or subconscious?

470

Psychology 402 - Spring 2020 - Dr. Michael Diehr

Internal vs. External Validity

- Internal Validity - how did it work?
 - were the methods good
 - did the IV cause the DV
- External Validity - what does it mean?
 - does skull size indicate IQ?
 - does IQ indicate personal worth?

471

Psychology 402 - Spring 2020 - Dr. Michael Diehr

Is skull size related to IQ?

- Yes. No. Maybe?
- Correlation between IQ and brain size: $R = 0.24$ (Pietschnig, 2015)
- $R^2 = \sim 6\%$ which means 94% of variance is *not explained*
- Thus, observed 3-4 cubic inch difference between the races would account for, *at most*, a 2-3 point IQ difference.
- But... men have ~10% larger brains than women, but do not show higher IQ (more on this later)

474

Psychology 402 - Spring 2020 - Dr. Michael Diehr

Review

- Frequency Distribution aka Histogram
- Normal Curve
 - CLT, Mean (SD)
- Law of Large Numbers
- Scales of Measurement
- Population vs. Sample
- Logical Fallacies
 - Precision vs. Accuracy
 - Reliability vs. Validity
 - Gambler's Fallacy

478

Psychology 402 - Spring 2020 - Dr. Michael Diehr

Comparing Scores

- Compare a single score to the population
- One way: difference scores
- Problem: Is a difference of “3” big or little? On a 100 point test it’s not very large, but on a 10 point test it’s the difference between an A and a C

479

Psychology 402 - Spring 2020 - Dr. Michael Diehr

Comparing Scores

- Desire a system independent of the raw score units (just like letter grades)
- Two methods:
 - Ranks & Percentile Ranks...
 - Standard Scores...

480

Psychology 402 - Spring 2020 - Dr. Michael Diehr

Ranks, Percentiles

- Given a distribution of scores, and a single score
- **Rank** = the item # of the single score when sorted high to low
- **Percentile Rank** = the % of scores which are lower than the given score
- **Percentile** = the score at which a given percent of scores are below a given score
- Note: “**Percentile**” often used informally to mean “**Percentile Rank**”

481

Psychology 402 - Spring 2020 - Dr. Michael Diehr

Rank & Percentile

- Infant mortality per 1000 live births
- Sorted low to high

Country	Score
Sweden	2.4
Japan	3.4
France	4.5
USA	7.5
Colombia	20.4
China	37.9
Bolivia	66.4
Ethiopia	142.6
Mozambique	148.6
Zambia	168.1

482

Psychology 402 - Spring 2020 - Dr. Michael Diehr

Rank & Percentile

- Determine Rank #

Country	Score	Rank
Sweden	2.4	1
Japan	3.4	2
France	4.5	3
USA	7.5	4
Colombia	20.4	5
China	37.9	6
Bolivia	66.4	7
Ethiopia	142.6	8
Mozambique	148.6	9
Zambia	168.1	10

483

Psychology 402 - Spring 2020 - Dr. Michael Diehr

Percentile Rank

Percentile Rank = # of cases with worse value divided by # of cases
e.g. France is third of 10 (it has 7 cases with worse values)
 $7 / 10 = 70\%$

Country	Score	Rank	%ile Rank
Sweden	2.4	1	90
Japan	3.4	2	80
France	4.5	3	70
USA	7.5	4	60
Colombia	20.4	5	50
China	37.9	6	40
Bolivia	66.4	7	30
Ethiopia	142.6	8	20
Mozambique	148.6	9	10
Zambia	168.1	10	0


Psychology 402 - Spring 2020 - Dr. Michael Diehr

484

Psychology 402 - Spring 2020 - Dr. Michael Diehr

485

Standard Scores



Psychology 402 - Spring 2020 - Dr. Michael Diehr

490

Standard Scores 2

- Use the mean and standard deviation as points of reference.
- Standard score : distance from the mean, scaled by standard deviation
- Not affected by raw score units.
- Different standard scores mean the same thing, but are expressed differently.
 - just like how 1.0 and 100% mean the same thing
- Unfortunately, there are several different Standard Score systems!

Psychology 402 - Spring 2020 - Dr. Michael Diehr

485

Standard Scores: Z, T, IQ

	Z scores	T scores	IQ scores
Mean	0	50	100
SD	1	10	15
Example: top 3%			
Example: top 1%			
Formula: from Z Score	Z	$(Z*10)+50$	$(Z*15)+100$

Psychology 402 - Spring 2020 - Dr. Michael Diehr

491

Norms 1

- Standard Scores provide us with a way of describing how a particular score relates to others in the population.
- Describing how an individual score relates to the population, which we assume are "normal".
- Terms "normative data" and "norms"
- Key questions: What is the normative group? What features or factors of the group may affect scores?

Psychology 402 - Spring 2020 - Dr. Michael Diehr

493

z-Score ($x - \bar{x}$) / s	T-Score $10z + 50$	Wechsler IQ ($15z + 100$)	Percentile Rank
3.0	80	145	99.9
2.9	79	144	99.8
2.8	78	142	99.7
2.7	77	141	99.6
2.6	76	139	99.5
2.5	75	138	99.4
2.4	74	136	99.2
2.3	73	135	98.9
2.2	72	133	98.6
2.1	71	132	98.2
2.0	70	130	97.7
1.9	69	129	97.1

Psychology 402 - Spring 2020 - Dr. Michael Diehr

492

Norms 2

- “norm-referenced” tests vs. “criterion-referenced” tests.
Example: an 85 year old in excellent shape could be in the top 5% of his class for firefighting ability, but this may still be a “failing” grade.
- Common factors that may matter:
- Gender, Age, Education, Ethnicity/Race, Language, Handedness, Height, Weight...