

# Ch. 2 - Measurement & Stats

122

Psychology 402 - Fall 2021 - Dr. Michael Diehr

Copyright © 2021 Michael Diehr  
All Rights Reserved  
For use only by students enrolled  
in my sections of Psyc 402  
through the end of the semester.  
May not be posted, shared or uploaded  
online without permission.

123

Psychology 402 - Fall 2021 - Dr. Michael Diehr

$$S = \sqrt{\frac{\sum_{i=1}^N (x_i - \bar{x})^2}{N - 1}}$$

126

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## Measurement & Stats

- Why numbers?
- Distribution & Graphs : Histogram
- Central Tendency
- Mean, SoR, SSR, Variance, Standard Deviation
- In-class exercise
- Population vs. Sample
- Measurement Scales
- Precision vs. Accuracy
- Logic and Logical Fallacies Descriptive vs. Inferential Statistics
- Norms

127

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## Basic Statistics

- Why use numbers?
- Pros:
  - convenient, succinct
  - universal
  - well-defined, repeatable
- Cons:
  - precision vs. accuracy
  - numerical fallacy

128


Psychology 402 - Fall 2021 - Dr. Michael Diehr

## Tabular Data

Year	Rainfall (inches)
1965	8.81
1966	14.76
1967	10.86
1968	7.86
1969	11.48

Psychology 402 - Fall 2021 - Dr. Michael Diehr

# Data Distributions



# Histogram


- Frequency Distribution
- Invented by Karl Pearson
- Shows data from *one* variable only
- Data is (often) collected into groups (“bins”)

131

Psychology 402 - Fall 2021 - Dr. Michael Diehr

# Histogram

Frequency Distribution of San Diego Annual Rainfall 1965-2017



# Describing Distributions

- Why? Large lists are inconvenient. Reduce many data points to a few numbers.
- Issue: Reducing data (“Degrees of freedom”): throws away data.
- We are modeling our data using a simplification.
- “All models are wrong, some models are useful”
- Simple vs. Simplistic?

133

Psychology 402 - Fall 2021 - Dr. Michael Diehr

# Descriptive Statistics

- Statistical Assumptions: When these are not met, weird things happen.
- Joe Smith is 6 feet tall, his child is 1 foot tall. Thus, the average height in the Smith household is 3.5 feet.
- If you are sitting in bar, and Bill Gates walks in, suddenly everyone in the bar is (on average) a multi-millionaire.

134

Psychology 402 - Fall 2021 - Dr. Michael Diehr

# Alternative Notations

- Square Root ( $x$ )  $\sqrt{x}$   $\sqrt[2]{x}$
- X-Squared  $X^2$   $X^{**2}$   $X^2$
- Sum( $x$ )  $x_1+x_2+x_3\dots$   $\sum_{i=1}^N x_i$   $\sum x$
- Mean  $M$   $\frac{\sum x}{N}$   $\bar{X}$

135

Psychology 402 - Fall 2021 - Dr. Michael Diehr

# Central Tendencies


- Values tend to cluster around a point.
- **Mean** : most common statistic, commonly referred to as the “average”. Formula  $\Sigma X / N$
- **Mode**: the most common value in a set
  - rare to use in statistics
- **Median**: the middle-most value in a set
  - the value at which half are above and half are below. Aka the 50th percentile.

136

Psychology 402 - Fall 2021 - Dr. Michael Diehr


# Histogram

Figure 1:  
Frequency Distribution of San Diego Annual Rainfall



Psychology 402 - Fall 2021 - Dr. Michael Diehr

# Normal Distribution




In a normal distribution, the mean, mode, and median are all the same

138

Psychology 402 - Fall 2021 - Dr. Michael Diehr

# Skewed Distribution




In a skewed distribution, the mean, mode, and median are all often different

139

Psychology 402 - Fall 2021 - Dr. Michael Diehr

# Skew

- negative skew : fatter tail on the left
- positive skew : fatter tail on the right



140

Psychology 402 - Fall 2021 - Dr. Michael Diehr

# Measures of Central Tendency 1

	Description	Algorithm	Formula
<b>Mean</b>	the “average”	sum values, divide by N	$\bar{x} = \frac{\sum_{i=1}^N x_i}{N}$
<b>Median</b>	the “middle-most value”	sort values, find middle value	50th percentile
<b>Mode</b>	the “most common” value	find most frequent value	...

141

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## Measures of Central Tendency 2

Behavior:	Normal Distribution	Skewed Distribution
<b>Mean</b>	same	overly affected by outliers
<b>Median</b>	same	fairly resistant to outliers
<b>Mode</b>	same	resistant to outliers

142

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## Measures of Dispersion 1

- Compare each measured value to the average
- “for a typical value, how far away is it from the mean”
- “Difference score” or “residual” can be calculated as the difference between the actual score and the mean. In other words,  $d_i = x_i - \bar{X}$
- Take the average (mean) of the difference scores.
- Average difference score =  $\text{Sum}(d) / N$

143

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## Average Difference Score

	Score (x)	Mean ( $\bar{X}$ )	Difference $d = (x - \bar{X})$
	2	6	-4
	3	6	-3
	9	6	3
	11	6	5
	14	6	8
	1	6	-5
	6	6	0
	4	6	-2
	5	6	-1
	5	6	-1
<b>Sum</b>	<b>60</b>	<b>60</b>	<b>0</b>
<b>Mean</b>	<b>6</b>	<b>6</b>	<b>0</b>

Psychology 402 - Fall 2021 - Dr. Michael Diehr

144

## Sum of Residuals

- Given N samples of x :  $x_1, x_2, x_3 \dots x_N$
- mean of x 
$$\bar{x} = \frac{\sum_{i=1}^N x_i}{N}$$
- residuals 
$$d_i = x_i - \bar{x}$$
- Sum of Residuals is always zero

$$\sum_{i=1}^N d_i = 0$$

145

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## Sum of Residuals

- The “average difference score” score will *always* equal zero
- Solution:
  - Square the residuals *before* adding: removes the negative values.
  - “SSR” or Sum of Squared Residuals

146

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## SSR: Sum of Squared Residuals<sup>147</sup>

- Given N samples of X:  $x_1, x_2, x_3 \dots x_N$
- mean of x 
$$\bar{x} = \frac{\sum_{i=1}^N x_i}{N}$$
- residuals 
$$d_i = x_i - \bar{x}$$
- Sum of Squared Residuals (SSR)

$$SSR = \sum_{i=1}^N (d_i)^2$$

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## Problems with SSR

- SSR depends on units of measurement:
  - a meter is 1000 millimeters, so SSR will be  $1000 \times 1000 =$  one million times higher when using meters vs. millimeters
- SSR depends on N (# of samples)
  - Doubling N will cause SSR to double (roughly)
- Therefore, SSR is hard to understand:
  - is  $SSR = 0.00342$  high or low?
  - is  $SSR = 2343249$  high or low?

148

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## Remove the influence of N

- The Sum of a set of values depends on the number (N) of values:
  - $\sum_{i=1}^N x_i$
- Take the average (mean)
  - this divides by N
  - removes the influence of N

149

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## Variance

- Problem: SSR depends on N
- Solution: Take the average of SSR to remove the influence of N
- The average of the squared residuals is called Variance ( $S^2$ )

150

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## Variance

- Variance =  $SSR/N$
- Variance = mean of squared residuals

$$S^2 = \frac{\sum_{i=1}^N (d_i)^2}{N}$$

$$S^2 = \frac{\sum_{i=1}^N (x_i - \bar{x})^2}{N}$$

151

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## Problems with Variance

- Units are squared:
  - measuring height in meters? variance is meters<sup>2</sup>
  - measuring # of cupcakes eaten? variance is (# of cupcakes eaten)<sup>2</sup>
- Won't someone rid me of these meddlesome squared units?

152

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## Standard Deviation

- Improving on Variance:
- The square root of Variance ( $S^2$ ) gives  $S$ , which is called "Standard Deviation".
- Also abbreviated SD, StdDev or  $\sigma$  (Greek letter sigma)
- SD : easier to understand because it's in the same units as your measurement.
- SD is a unique property of the normal distribution -- given a mean and a SD you have uniquely specified the distribution

153

Psychology 402 - Fall 2021 - Dr. Michael Diehr

# Standard Deviation

- SD = Square root of Variance

$$S = \sqrt{\frac{\sum_{i=1}^N (d_i)^2}{N}}$$

$$S = \sqrt{\frac{\sum_{i=1}^N (x_i - \bar{x})^2}{N}}$$

154

Psychology 402 - Fall 2021 - Dr. Michael Diehr


# Standard Deviation

- can be thought of as the “average deviation”
- (but it’s not literally average deviation, since we showed earlier the average difference score is always Zero)
- Technically:
  - (in a normal distribution) scores will be within plus or minus 1 SD about 68% of the time

155

Psychology 402 - Fall 2021 - Dr. Michael Diehr

# Normal Distribution



In a normal distribution, about 68.2% of values fall within  $\pm 1$  SD


156

Psychology 402 - Fall 2021 - Dr. Michael Diehr

# SSR, Variance and SD

157

Figure 1: Frequency Distribution of San Diego Annual Rainfall



Psychology 402 - Fall 2021 - Dr. Michael Diehr

# Central Limit Theorem

- No matter the shape of the Population distribution, if you take enough (\*) samples of the mean, the distribution of your samples of the mean will have a Normal distribution
- Central Limit Theorem Exercise (Javascript)
- This fact makes our life easy: Many statistics assume a normal distribution. The CLT provides us a normal distribution in most cases, even when the population data is skewed

158

Psychology 402 - Fall 2021 - Dr. Michael Diehr

# Exercise: normal distribution

- Roll one 10-sided die 10 times and record the results
- Prediction
  - Your Distribution: Uniform (flat)
  - Mean : 4.5
  - Class Distribution: ???
- hint: What is N? # die rolls, # of students?
- List and Graph results
- Does the distribution look normal?
  - if so, why?

159

Psychology 402 - Fall 2021 - Dr. Michael Diehr

### Exercise 1: Die Rolls

X	M	$d = (x_i - M)$	$(\text{residual})^2$
2	3	-1	1
3	3		
5	3		
2	3		


N	$M = \bar{X}$	$\Sigma \text{Residuals}$	$\Sigma (\text{residual}^2)$	$\frac{\Sigma (\text{residual}^2)}{N-1}$	$\sqrt{S^2}$
	3				

160

Psychology 402 - Fall 2021 - Dr. Michael Diehr

Title:

**Figure 1**  
Blah de blah blah blah



Y axis label:

\_\_\_\_\_

X axis label:

Note:


161

Psychology 402 - Fall 2021 - Dr. Michael Diehr

### Example of APA-7 style Histogram

**Figure 1**

Frequency Distribution of annual Rainfall in San Diego



Psychology 402 - Fall 2021 - Dr. Michael Diehr

162

### Exercise: normal distribution 2

- Compute Mean ( $\bar{X}$ ) - is it near 4.5?
- Compute residuals
- Compute sum of residuals -- do they add to zero?
- Compute squared residuals
- Compute Sum of squared residuals (SSR)
- Divide SSR by  $(N-1)$  - this is Variance or  $(S^2)$
- Take square root of variance - this is S or Standard Deviation
- For this exercise, SD should be near 2.8

163

Psychology 402 - Fall 2021 - Dr. Michael Diehr

### Class Data: 1pm section

Mean of Die Rolls	# Students
0	
1	
2	
3	
4	
5	
6	
7	
8	
9	

164

Psychology 402 - Fall 2021 - Dr. Michael Diehr

### Class Data: 4pm section

Mean of Die Rolls	# Students
0	
1	
2	
3	
4	
5	
6	
7	
8	
9	

165

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## Ch. 2 - Part 2

167

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## Review - Stats

- Why Numbers?

- pros
- cons

- Distributions

- Tables
- Graphs

169

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## Review - Descriptives

- Frequency Distribution aka Histogram
  - graphically shows data
- Central Tendency
  - mean, median, mode
- Dispersion or Variation
  - residual
  - sum of residuals = 0
  - sum of squared residuals > 0
  - SSR/N = Variance
  - $\text{Sqrt}(\text{Variance})$  = Standard Deviation

170

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## Frequency Distributions


171

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## SSR, Variance and SD


Figure 1:

Frequency Distribution of San Diego Annual Rainfall



172

## Review : Distributions



In a normal distribution, about 68.2% of values fall within  $\pm 1$  SD

173

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## Standard Deviation

- SD = Square root of Variance

$$S = \sqrt{\frac{\sum_{i=1}^N (d_i)^2}{N}}$$

$$S = \sqrt{\frac{\sum_{i=1}^N (x_i - \bar{x})^2}{N}}$$

174

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## Exercise: key points

- Some events (such as the roll of a die) have a flat (or ‘uniform’) distribution, but these are rare.
- Many big events are composed of many small events.
- Events in the real world often are distributed in a (nearly) normal distribution
- Assuming a normal distribution, the easiest way to describe the data is by two factors: Mean and SD.

175

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## Central Limit Theorem

- No matter the shape of the Population distribution, if you take enough (\*) samples of the mean, the distribution of your samples of the mean will have a Normal distribution
- [Central Limit Theorem Exercise \(Javascript\)](#)
- This fact makes our life easy: Many statistics assume a normal distribution. The CLT provides us a normal distribution in most cases, even when the population data is skewed

176

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## Types of Statistics

- Descriptive:
  - Goal: help you describe the data
  - reduce the amount of data necessary for understanding
  - don’t draw conclusions -- “just the facts”
- Inferential:
  - Goal: draw conclusions from your sample to the larger data set (population)

177

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## Measurement Scales

- Nominal
- Ordinal
- Interval
- Ratio

178

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## Nominal Scale

- Nominal: Name or ID only
  - red, blue, green....
  - john, tony, fred...
  - Sci2-243, Sci2-245...
- does not signify Ordering, Ranking, or More/Less
- Gotcha: even if used with Numbers it may be still a Nominal.
- Example: colors, names, room numbers, ID numbers

179

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## Ordinal Scale

- Ordinal : ordering
- first, second, third....  
1, 2, 3...  
A, B, C...
- signifies Order, but can't assume distance between items is the same, e.g. the difference between an A and a B may be much different than a B and a C
- Example: Class Rank, Assignment Grade, Product Ratings

180

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## Interval Scale

- Interval: specifies orders AND inter-item distance
  - -3, -2, -1, 0, 1, 2, 3.... 100, 105, 115
  - the difference between two numbers IS the same, e.g. 100 to 105 should be the same amount as 105 to 110
- Does NOT have an absolute zero.
- Example: temperature in Degrees Farenheit

181

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## Ratio Scale

- Ratio: specifies orders AND inter-item distance and has absolute zero
- 0, 1, 2, 3.... 100, 105, 115
- the difference between two numbers IS the same, e.g. 100 to 105 should be the same amount as 105 to 110
- Does have an absolute zero.
- Example: temperature in Degrees Kelvin

182

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## Measurement Scales

	Magnitude	Equal Intervals	Absolute Zero
Nominal			
Ordinal	✓		
Interval	✓	✓	
Ratio	✓	✓	✓

183

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## Scales: Practical Info

- Nominal Scale: common
  - common stats: Count, Frequency, Mode
- Ordinal Scale: less common
  - stats: specialized “nonparametric” techniques required
- Ratio and Interval: common
  - Often can be treated identically with same statistical techniques

184

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## Descriptive Statistics

- Count (N)
- Range (minimum, maximum)
- Frequency Distribution (histogram)
- Rank order, percentile (%ile)
- Central Tendency
  - Mean
  - Median
  - Mode
- Variation / Dispersion (Variance, Standard Deviation)

185

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## Population vs. Sample

- Ideally, measure *everyone* to get the exact value (*Population parameter*)
- Practically, this is impossible.
- Take samples instead, and calculate the *Sample statistic*.
- The “Law of Large Numbers”, “Sampling Theory”, “Central Limit Theorem” makes life easier
- Central Limit Theorem Exercise (Javascript)
- Some formulas differ for *Population vs. Sample* (divide by N or divide by N-1 ?)

186

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## Population v. Sample

	Population	Sample
<b>Definition</b>	the entire set of items	the actual subset you measured
<b>Descriptives</b>	“Parameters”	“Statistics”
<b>Symbols</b>	Greek	Roman
<b>Mean</b>	$\mu$	$\bar{x}$
<b>Std. Deviation</b>	$\sigma$	$s$
<b>Variance</b>	$\sigma^2$	$s^2$
<b>Divide by</b>	N	N-1

187

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## Law of Large Numbers

- If you take enough\* samples, the sample mean approaches the population mean.
- Example: a coin has two sides. If heads=1 and tails = 0, then the average expected result is exactly 50% Heads (0.5) in the long run.
- However, if you flip a coin just a few times, getting exactly 0.5 is not likely.
- The LLN states that you will if you take enough samples.

\* what is “enough”? Rule of thumb : 100.

188

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## LLN Demonstration

- Law of Large Numbers
- Demonstration with Coin Flips

189

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## Logical Fallacies

190

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## Logical Arguments

- Logical arguments or inferences generally have several components:
  - Premises
  - Conclusions
- Example:
  - Premise: All English people are musicians
  - Premise: John Lennon was English
  - Conclusion: John Lennon was a musician

192

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## Logical Arguments 2

- An Inference can be either Valid or Invalid -- this refers to the Structure of the argument (not the Facts themselves)
  - All A are B
  - All C are A
  - All C are B
- A Valid inference can still come to a false conclusion, and vice-versa

193

Psychology 402 - Fall 2021 - Dr. Michael Diehr


## Logical Fallacies

- A Logical Fallacy generally means that your inference is Invalid to begin with. In addition, your facts may or may not be true, but the flaw in reasoning has occurred before you even apply facts.
  - Example: Affirming the consequent
  - If P, then Q
  - Q is true
  - Therefore P
- bank owners are rich
- Bill Gates is rich
- Bill Gates works at a bank

194

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## Cognitive Bias Codex



196

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## Biased Sample

- Every individual  $x$  that we have seen from sample X has characteristic Z  
Therefore ALL X have characteristic Z
- Every student I talk to in this class is interested in Psychology  
Therefore, ALL students are interested in Psychology

## Nominal and Numeric

- Nominal Fallacy:** The tendency to believe that something has a name or identification, it exists or has special meaning.  
“I am sleepy” vs. “I am suffering from activity-induced-rest-reduction-performance-impairment syndrome”
- Numerical Fallacy:** belief that something has been measured and assigned a number, it actually exists. “I’m really sad” vs. “I scored a 32 on the Beck Depression Inventory”

197

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## Reification Fallacy

- To Reify - to make something more concrete or real
- Examples:
  - “An A student”
  - “High IQ”
  - “Top of the class”
  - “A F Grade”

198

Psychology 402 - Fall 2021 - Dr. Michael Diehr


## Ranking Fallacy

- Reducing a complex phenomenon (e.g. intelligence), giving it a single number (reification) and then ordering based on that number
- Examples:
  - A IQ of 93 is better than an IQ of 90
  - An income of \$50,000 is better than \$45,000

199


Psychology 402 - Fall 2021 - Dr. Michael Diehr

## Correlation = Causation



200

## Hasty Generalization



201

## Other Fallacies

- Begging the question -- circular argument
- Correlation implies causation
- Post hoc ergo propter hoc (*after this, therefore because of this*)
- Appeal to Authority
- Ad-hominem
- Straw Man
- False Dilemma

202

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## Significant Figures

- “digits of precision” or “sig. fig.”
- Ignoring *leading zeros*, how many digits in the measurement?
  - 00123 has 3 sig. fig.
  - 12003 has 5 sig fig
  - 9.87 has 3 sig fig
  - 0.000987 has 3 sig. fig.
  - 12.1 has 3 sig. fig.
  - 12.0 has 3 sig. fig.
- Please use 3 sig fig for this class
- Note: not the same as “3 decimal places”

203

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## Precision vs. Accuracy

- Precision : the level of detail a measurement is made with, often specified with an error-range
  - “about 6 feet plus or minus 1 foot” vs. “6 foot 11 inches plus or minus 1 inch”
- Accuracy: how close the measured value is to the actual value, does it “hit the target”
  - Think arrow vs. shotgun
- A number can be precise and accurate, precise but inaccurate, or accurate but imprecise.

204

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## Precision Fallacy


- A number that is *precise* may seem to be *accurate* when it is not
- A measurement that is *reliable* may seem to have *validity* when it does not

205

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## Precision vs. Accuracy

- Target shooting analogy
- Similar to Reliability vs. Validity



Psychology 402 - Fall 2021 - Dr. Michael Diehr

## Fallacies re: Probability

- Classical
- Gambler's Fallacy
- Bayesian Reasoning

207

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## 9 Heads in a row

- You are flipping a coin, and get 9 heads in a row  
H H H H H H H H H
- What is the % chance the next flip will be a H ?
- Three common answers:
  - 50/50
  - more likely Heads
  - more likely Tails

208

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## 9 Heads: Classical Inference

- Coin flips are independent 50/50 events, therefore 50% : Logical/Statistical
- This is the \*correct\* answer for a fair coin

209

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## LLN Demonstration

- Law of Large Numbers
- Demonstration with Coin Flips

210

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## 9 Heads: Gambler's Fallacy

- Coin flips are independent 50/50 events, but we just saw 9/10 heads, therefore a Tail is “due”
- This is the “Gambler’s Fallacy” and one reason Casinos make tons of money. The reasoning is faulty.
- Note: when dealing with draws w/o replacement, this logic is *\*correct\**. For example, a single-card blackjack deck -- if no face cards have come up after 30 cards, then face cards are due

211

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## 9 Heads: Bayesian Statistics

- Coin flips are supposed to be 50/50 events, but we just saw 9/10 heads, therefore the data is telling us that perhaps this is not a fair coin.
- Bayes’ theorem suggests you evaluate the prior probabilities in determining future behavior
- In this case, you’d conclude that Head is more likely on the next flip

212

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## Cognitive Biases of Discrimination

- Which cognitive biases (logical fallacies) are involved in racism, sexism and other bigoted beliefs?

213

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## Ch. 2 - Part 3

## Review - History


- Themes
  - 18th-19th century
  - 19th-20th century
- Theories of Human Development
  - Creationism
  - Polygenism
  - Evolution
  - Genetics
- Controversy
  - IQ testing of various groups

217

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## Louis Agassiz

- Swiss-born, European-trained biologist / geologist
- Came to Harvard in 1847
- Creationist -> Polygenist
- Taxonomist
- Resisted Darwin’s theory of Evolution
- d. 1873




218

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## Review: Samuel George Morton


- Theory of Polygenism
- Humans are composed of different species, created by god
- Craniometry
- Biological Determinism
- “Scientific Racism”
- The “American School”
- d. 1851



219

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## Crania Americana



Samuel George Morton  
1839

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## Morton's Data as printed

Race	N	Cranial Volume Mean
Caucasian	52	87
Mongolian	10	83
American	144	82
Malay	18	81
Ethiopian	29	78

223

Psychology 402 - Fall 2021 - Dr. Michael Diehr


## Data, corrected

Race	Mean (Morton)	Mean (corrected)
Caucasian	87	87
Mongolian	83	87
American	82	86
Malay	81	85
Ethiopian	78	83

224

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## As Printed



225

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## Seed vs. Shot

Race	Difference (seed - shot)
Caucasian	1.8
Mongolian	n/a
American	2.2
Malay	n/a
Ethiopian	5.4

226

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## Morton's errors

- Fundamental arithmetic errors
- Data selection errors
- Failure to measure or control for external variables (biological sex, body size, etc.)
- Basic Statistical errors (averaging measurements from unequal size subgroups)
- The racist thumb press?
- Is he a liar? Conscious or subconscious?

227

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## Morton's Model

228

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## Internal vs. External Validity

- Internal Validity - how did it work?
  - were the methods good
  - did the IV cause the DV
- External Validity - what does it mean?
  - does skull size indicate IQ?
  - does IQ indicate personal worth?

229

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## Is skull size related to IQ?

- Yes. No. Maybe? Probably not much.
- Correlation between IQ and brain size:  $R = 0.24$  (Pietschnig, 2015)
- $R^2 = \sim 6\%$  which means 94% of variance is *not explained*
- Thus, observed 3-4 cubic inch difference between the races would account for, *at most*, a 2-3 point IQ difference.
- But... men have  $\sim 10\%$  larger brains than women, but do not show higher IQ (more on this later)

230

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## Comparing Scores

- Compare a single score to the population
- One way: difference scores
- Problem: Is a difference of "3" big or little? On a 100 point test it's not very large, but on a 10 point test it's the difference between an A and a C

231

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## Comparing Scores

- Desire a system independent of the raw score units (just like letter grades)
- Two methods:
  - Ranks & Percentile Ranks...
  - Standard Scores...

232

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## Ranks, Percentiles

- Given a distribution of scores, and a single score
- Rank** = the item # of the single score when sorted high to low
- Percentile Rank** = the % of scores which are lower than the given score
- Percentile** = the score at which a given percent of scores are below a given score
- Note: “Percentile” often used informally to mean “Percentile Rank”

233

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## Rank & Percentile

- Infant mortality per 1000 live births
- Sorted low to high

Old slide -  
From textbook  
please ignore

Country	Score
Sweden	2.4
Japan	3.4
France	4.5
USA	7.5
Colombia	20.4
China	37.9
Bolivia	66.4
Ethiopia	142.6
Mozambique	148.6
Zambia	168.1

234

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## Rank & Percentile

- Coronavirus Deaths
- Total deaths, per million people
- Sort low to high

Country	Score
Mozambique	0.9
China	3.0
Ethiopia	8.0
Japan	11.0
Zambia	16.0
Colombia	424.0
France	471.0
Sweden	577.0
USA	584.0
Bolivia	599.0

235

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## Rank & Percentile

- Determine Rank #

Country	Score	Rank
Mozambique	0.9	1
China	3.0	2
Ethiopia	8.0	3
Japan	11.0	4
Zambia	16.0	5
Colombia	424.0	6
France	471.0	7
Sweden	577.0	8
USA	584.0	9
Bolivia	599.0	10

236

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## Percentile Rank

Percentile Rank = # of cases with worse value divided by # of cases

e.g. France is 7th of 10 (it has 3 cases with worse values)  
 $3 / 10 = 30\%$  percentile rank

Country	Score	Rank	%ile Rank
Mozambique	0.9	1	90
China	3.0	2	80
Ethiopia	8.0	3	70
Japan	11.0	4	60
Zambia	16.0	5	50
Colombia	424.0	6	40
France	471.0	7	30
Sweden	577.0	8	20
USA	584.0	9	10
Bolivia	599.0	10	0

237

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## Standard Scores 2

- Use the mean and standard deviation as points of reference.
- Standard score : distance from the mean, scaled by standard deviation
- Not affected by raw score units.
- Different standard scores mean the same thing, but are expressed differently.
  - just like how 1.0 and 100% mean the same thing
- Unfortunately, there are several different Standard Score systems!

238

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## Z-score


- A Z score is the # of standard deviations above (+) or below (-) the mean of a single measurement.
- Algorithm: given a single score ( $X_i$ ), subtract the mean  $M$ , divide by the standard deviation  $S$
- Formula
  - $Z = (X-M)/SD$

$$Z_i = \frac{X_i - \bar{X}}{S}$$

239

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## Standard Scores



## Standard Scores: Z, T, IQ

	Z scores	T scores	IQ scores
<b>Mean</b>	0	50	100
<b>SD</b>	1	10	15
<b>Example: top 3%</b>			
<b>Example: top 1%</b>			
<b>Formula: from Z Score</b>	$Z$	$(Z*10)+50$	$(Z*15)+100$

241

Psychology 402 - Fall 2021 - Dr. Michael Diehr

z-Score ( $X-\bar{X})/S$	T-Score $10z + 50$	Wechsler IQ $(15z + 100)$	Percentile Rank
3.0	80	145	99.9
2.9	79	144	99.8
2.8	78	142	99.7
2.7	77	141	99.6
2.6	76	139	99.5
2.5	75	138	99.4
2.4	74	136	99.2
2.3	73	135	98.9
2.2	72	133	98.6
2.1	71	132	98.2
2.0	70	130	97.7
1.9	69	129	97.1

242

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## Norms 1

- Standard Scores provide us with a way of describing how a particular score relates to others in the population.
- Describing how an individual score relates to the population, which we assume are “normal”.
- Terms “normative data” and “norms”
- Key questions: What is the normative group? What features or factors of the group may affect scores?

243

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## Norms 2

- “norm-referenced” tests vs. “criterion-referenced” tests.  
Example: an 85 year old in excellent shape could be in the top 5% of his class for firefighting ability, but this may still be a “failing” grade.
- Common factors that may matter:
- Gender, Age, Education, Ethnicity/Race, Language, Handedness, Height, Weight...

244

Psychology 402 - Fall 2021 - Dr. Michael Diehr


## Z-score Exercise

- This is for practice, not graded for points
- PDF is on class website

245

Psychology 402 - Fall 2021 - Dr. Michael Diehr

## Standard Scores



## Standard Scores: Z, T, IQ

	Z scores	T scores	IQ scores
Mean	0	50	100
SD	1	10	15
Example: top 3%	1.9		
Example: top 1%	2.4		
Formula to convert from Z Score	Z	$(Z \times 10) + 50$	$(Z \times 15) + 100$

247

Psychology 402 - Fall 2021 - Dr. Michael Diehr